

Wisconsin Ag News – Crop Progress & Condition



Upper Midwest Regional Field Office \cdot 210 Walnut St, Ste 833 \cdot Des Moines, IA 50309 \cdot (515) 776-3400 www.nass.usda.gov/wi

Cooperating with the Wisconsin Department of Agriculture, Trade and Consumer Protection

June 30, 2025 - For Immediate Release

Wisconsin had 3.3 days suitable for fieldwork statewide for the week ending June 29, 2025, according to the USDA's National Agricultural Statistics Service. Heavy rain was received across large portions of the state, limiting fieldwork.

Topsoil moisture condition rated 1 percent very short, 11 percent short, 67 percent adequate and 21 percent surplus. **Subsoil moisture** condition rated 3 percent very short, 12 percent short, 71 percent adequate and 14 percent surplus.

Nearly all the **corn** crop had emerged. Corn condition was rated 75 percent good to excellent, 1 percentage point below last week. **Soybeans** were 97 percent emerged. Ten percent of soybeans were blooming, matching both last year and the 5-year average. Soybean condition was 73 percent good to excellent, unchanged from last week.


Winter wheat was 95 percent headed. Forty-one percent had begun coloring, 5 days behind last year, and 4 days behind the average. Condition was rated 72 percent good to excellent, down 3 percentage points from last week. Oats were 98 percent emerged. Oats were 63 percent headed, 4 days behind last year, and 3 days behind the average. Ten percent of oats were reported to have begun coloring. Oat condition was rated 85 percent good to excellent, down 2 percentage points from last week. Potato condition was 86 percent good to excellent, down 2 percentage points from last week.

The first cutting of **alfalfa hay** was 94 percent complete. The second cutting was 21 percent complete, 1 day ahead of last year, but 2 days behind the average. **Hay** condition rated 82 percent good to excellent, up 4 percentage points from last week. **Pasture and range** condition rated 74 percent good to excellent, 1 percentage point above last week.

Crop Condition as of June 29, 2025

Item	Very Poor	Poor	Fair	Good	Excellent	
	(percent)	(percent)	(percent)	(percent)	(percent)	
Corn	1 0 0 1	3 2 2 5 0 4	21 16 13 20 14 22	58 65 67 57 80 57	17 17 18 17 6	
Soybeans Wheat, winter	1	7	20	55	17	





Crop Progress as of June 29, 2025

Item	Districts									State			
	NW	NC	NE	WC	С	EC	SW	sc	SE	This week	Last week	Last year	5-year avg
	(percent)	(percent)	(percent)										
Hay, alfalfa, 1st cutting	94	66	99	95	89	98	97	99	98	94	90	89	96
Hay, alfalfa, 2nd cutting		0	25	18	8	35	18	39	48	21	10	20	26
Oats headed	57	43	79	61	36	39	93	80	79	63	47	72	69
Oats coloring	0	0	0	12	10	2	20	26	24	10	3	16	17
Soybeans emerged	98	91	99	99	91	97	98	100	98	97	94	95	97
Soybeans blooming	1	0	2	4	10	3	20	23	6	10	3	10	10
Wheat, winter, headed	87	71	94	92	82	96	100	99	99	95	91	98	95
Wheat, winter, coloring	5	2	11	21	16	38	64	57	63	41	15	57	52


Days Suitable for Fieldwork and Soil Moisture Condition as of June 29, 2025

			State									
Item	NW	NC	NE	WC	O	EC	SW	SC	SE	This week	Last week	Last year
	(days)	(days)										
Days suitable	3.3	0.5	3.9	2.5	3.6	4.6	2.0	4.3	5.5	3.3	4.8	3.3
	(percent)	(percent)										
Topsoil moisture												
Very short	0	0	0	0	0	0	0	0	16	1	1	0
Short	7	3	3	1	1	12	12	28	23	11	19	0
Adequate	61	33	69	83	73	72	74	65	59	67	70	48
Surplus	32	64	28	16	26	16	14	7	2	21	10	52
Subsoil moisture												
Very short	0	0	1	0	0	0	5	4	24	3	3	0
Short	3	1	4	1	1	9	20	32	22	12	15	0
Adequate	70	50	77	90	79	80	70	58	52	71	74	59
Surplus	27	49	18	9	20	11	5	6	2	14	8	41

The complete report can be found on the USDA NASS website at www.nass.usda.gov/Publications.


Average Temperature (°F): Departure from 1991-2020 Normals

June 23, 2025 to June 29, 2025



Accumulated Precipitation (in)

June 23, 2025 to June 29, 2025



Temperature and Precipitation Maps, courtesy of the Midwestern Regional Climate Center, are available at: https://mrcc.purdue.edu/CLIMATE/

Additional soil moisture data are available at: https://nassgeo.csiss.gmu.edu/CropCASMA/